

# Fenadin<sup>®</sup>

(Fexofenadine Hydrochloride USP)

## DESCRIPTION:

**Fenadin<sup>®</sup> 60** : Each tablet contains Fexofenadine Hydrochloride 60mg

**Fenadin<sup>®</sup> 120** : Each tablet contains Fexofenadine Hydrochloride 120mg

**Fenadin<sup>®</sup> 180** : Each tablet contains Fexofenadine Hydrochloride 180mg

## INDICATIONS & USAGE:

### Seasonal Allergic Rhinitis

**Fenadin<sup>®</sup>** is indicated for the relief of symptoms associated with seasonal allergic rhinitis in adults and children 2 years of age and older. Symptoms treated effectively include

- Sneezing
- Rhinorrhea
- Itchy & Watery nose and throat
- Itchy skin
- Red eyes

### Chronic Idiopathic Urticaria

**Fenadin<sup>®</sup>** is also indicated for the treatment of uncomplicated skin manifestations of chronic idiopathic urticaria in adults and children 6 months to 11 years of age.

## DOSAGE AND ADMINISTRATION:

### Fenadin<sup>®</sup> Tablet

#### Seasonal Allergic Rhinitis

**Adults and Children 12 years and older** : The recommended dose of **Fenadin<sup>®</sup>** is 60mg twice daily or 120mg or 180mg once daily. A dose of 60mg once daily is recommended as the starting dose in patients with decreased renal function.

**Children 6 to 11 years** : The recommended dose of **Fenadin<sup>®</sup>** is 30mg twice daily. A dose of 30mg once daily is recommended as the starting dose in pediatric patients with decreased renal function.

### Chronic Idiopathic Urticaria

**Adults and Children 12 years and older** : The recommended dose of **Fenadin<sup>®</sup>** is 60mg twice daily. A dose of 60mg once daily is recommended as the starting dose in patients with decreased renal function.

**Children 6 to 11 years** : The recommended dose of **Fenadin<sup>®</sup>** is 30mg twice daily. A dose of 30mg once daily is recommended as the starting dose in pediatric patients with decreased renal function.

## SIDE EFFECTS:

The incidence of adverse events, including drowsiness, is not dose related and is similar across subgroups defined by age, gender and race. The other side effects are Viral like infection (cold, flu), nausea, dysmenorrhea, fatigue, headache and throat irritation.

## DRUG INTERACTION:

**Fenadin<sup>®</sup>** should not be taken closely in time with Aluminium and Magnesium containing Antacid, Erythromycin or Ketoconazole co-administration may enhance Fexofenadine absorption.

## CONTRAINDICATION:

Fexofenadine Hydrochloride is contraindicated in patients with known hypersensitivity to any of its ingredients.

## PREGNANCY:

There are no adequate and well-controlled studies in pregnant women. **Fenadin<sup>®</sup>** should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

## NURSING MOTHERS:

No adverse effects have been seen in breast feeding infants whose mothers were receiving Fexofenadine Hydrochloride and **THE AMERICAN ACADEMY OF PEDIATRICS** considers that it is therefore usually compatible with breast feeding.

## PEDIATRIC USE:

Safety and effectiveness of Fexofenadine Hydrochloride in pediatric patients can be used up to the age of 6 months.

## STORAGE:

Store in a cool and dry place, away from light & children.

## PACKAGE QUANTITIES:

**Fenadin<sup>®</sup> 60** : Each box containing 3X10's film coated tablets in blister strips.

**Fenadin<sup>®</sup> 120** : Each box containing 1X10's, 2X10's, 5X10's film coated tablets in blister strips.

**Fenadin<sup>®</sup> 180** : Each box containing 2X10's film coated tablets in blister strips.

<sup>®</sup> Trade Mark

Manufactured by  
 **Renata Limited**  
Mirpur, Dhaka, Bangladesh

Tablet (60mg, 120mg & 180mg)

Manufactured by  
 **Renata Limited**  
Rajendrapur, Gazipur, Bangladesh.

Tablet (60mg, 120mg & 180mg)