

Domiren®

Domperidone Maleate BP

Description :

Domiren® is a dopamine antagonist that principally blocks the dopamine receptors located in the CTZ and stomach. **Domiren®** restores normal motility and tone of the upper gastrointestinal tract, facilitates gastric emptying, enhances antral and duodenal peristalsis and regulates contraction of the pylorus. **Domiren®** has almost no effect on the dopaminergic receptors in the brain, therefore, excludes psychotropic and neurologic side effects.

Composition :

Domiren® Tablet : Each tablet contains Domperidone maleate BP equivalent to 10mg of Domperidone.

Domiren® Paediatric Drops : Each ml contains Domperidone BP 5mg.

Domiren® Suspension : Each 5ml suspension contains Domperidone BP 5mg.

Indications :

- Stimulation of gut mobility in non-ulcer dyspepsia, esophageal reflux, reflux esophagitis and gastritis, diabetic gastroparesis, functional dyspepsia.
- Prevention and symptomatic relief of acute nausea and vomiting from any cause including cytotoxic therapy, radiotherapy, antiparkinsonism therapy.
- Treatment of acute migraine.
- Correction of defective and insufficient lactation.

Dosage and administration :

The recommended oral dose for

Adults : 10-20mg (1-2 tablets or 10-20ml suspension) every 4-8 hours daily.

Children : 2-4ml suspension /10kg Or 0.4-0.8ml Drops/10 kg body weight every 4-8 hours daily.

For acute nausea and vomiting, maximum period of treatment is 12 weeks

Note : **Domiren®** should be taken 15-30 minutes before meals.

Side effects :

Domperidone may produce hyperprolactinemia (1.3% frequency). This may result in galactorrhea, breast enlargement and soreness and reduced libido. Dry mouth (1.9%), thirst, headache (1.2%), nervousness, drowsiness (0.4%), diarrhea (0.2%), skin rash and itching (0.1%) may occur during treatment with Domperidone. Extra-pyramidal reactions are seen in 0.05% of patients in clinical studies.

Contraindications :

Domperidone is contraindicated to the patients who have known hypersensitivity to this drug and in case of neonates.

Precautions :

Domperidone should be used with caution in case of children because there may be an increased risk of extra-pyramidal reactions in young children because of an incompletely developed blood brain barrier.

Use in pregnancy & lactation :

Pregnant women : The safety of Domperidone has not been proven and it is therefore not recommended in pregnancy.

Lactating women : Domperidone may precipitate galactorrhea and improve postnatal lactation. It is secreted in breastmilk but in very small quantities, insufficient to be considered harmful.

Drug interactions :

Domperidone may reduce the hypoprolactinemic effect of bromocriptine. The action of domperidone on GI function may be antagonized by anti-muscarinics and opioid analgesic.

Overdosage :

There is no reported case of overdosage.

Storage condition :

Store in a cool and dry place, away from light and children.

Commercial pack :

Domiren® Tablet : Each box contains 10X10's, 12X10's, 15X10's film coated tablets in Alu-PVC blister pack.

Domiren® Paediatric Drops : Each box contains 15ml of paediatric drops in a glass bottle.

Domiren® Suspension : Each box contains 60ml and 100ml of suspension in a glass bottle.



Manufactured by
Ziska Pharmaceuticals Ltd. for
RENATA LIMITED
Mirpur, Dhaka, Bangladesh
©Trademark